



STUDY ON RELATIONSHIP BETWEEN HEART RATE VARIABILITY BETWEEN DOSHA PRAKRUTI: AN OBSERVATIONAL STUDY

Amit Kumar Mishra^{1*}, Manisha V. Bhalsing², Sapna Upadhyay¹, Abdul Mutallib³ and Satish Bajpai¹

¹Assistant Professor, Department of Kriya Sharir, Babu Yugraj Singh Ayurvedic Medical College and Hospital, Gomti Nagar, Lucknow, UP, India.

²Professor, Department of Kriya Sharir, Bharati Vidyapeeth Deemed University, Pune, India.

³PG Scholar, Department of Preventive and Social Medicine, ZVM Unani Medical College, Pune, India.

Article Received on
24 January 2022,

Revised on 14 Feb. 2022,
Accepted on 04 March 2022

DOI: 10.20959/wjpps20224-16632

*Corresponding Author

Dr. Amit Kumar Mishra

Assistant Professor,
Department in Kriyasharir in
Babu Yugraj Singh
Ayurvedic Medical College
and Hospital Lucknow UP.
dramitmishra1989@gmail.com

ABSTRACT

Dosha Prakruti is an important physiological parameter of Ayurveda, and its derangement in various diseases is basis the pathological process of disorders. The scientific study about relationship between Dosha- Prakruti and heart rate variability is one of the most explored and relevant topics in the present scenario of dramatically increasing prevalence of cardiovascular diseases in the elderly population. The present observational study is an endeavor to explore the causal relationship between HRV and Dosha in the older adults and elderly populations and impact of exercise on these parameters. The study inferred that individuals with Vata Dosha are highly prone to develop increased HRV that underlines the propensity of developing sinus dysfunction and exercise is an important mode to stimulate the vagal tone for the healthy heart.

KEYWORDS: Dosha, HRV, Vata, Elderly.

INTRODUCTION

Ayurveda is the system of medicine with a rationale logical foundation which has survived as a distinct preventive and therapeutic entity from antiquity to the present day of sophisticated medical innovations.^[1,2] The modern scientific innovations have logically proved the basic tenets of Ayurveda in the maintenance and cure of human ailments. Dosha Prakruti is an

important physiological parameter of Ayurveda, and its derangement in various diseases is basis the pathological process of disorders.^[3] The scientific study about relationship between Dosha- Prakruti and heart rate variability is one of the most explored and relevant topics in the present scenario of dramatically increasing prevalence of cardiovascular diseases in the elderly population.^[4,5]

The human body is a biological combination of the five elements: Ether, Air, Fire, Water, and Earth. The combination of Ether and Air form Vata, Fire and Water form Pitta, and Water and Earth form Kapha^[6,7] The balance of these Dosha determines the maintenance of health, whereas the imbalance of Dosha may result in failure of hemodynamic stability and health.^[8]

Since ancient times, Ayurvedic physicians have used various methods of examination to determine Doshas^[9,10] such as observation of the body, body functions, reading of the pulse, examination of the eyes, memory, and behavioral habits such as religious observation, and sleep and eating habits.^[10]

The variations in Tridoshas are the basis for disease diagnosis and treatment in Ayurveda.^[11] The *Doshas* are assessed by sensing the pulse manually with fingers which depends on skill of the physician.^[3] Thus, imbalance in Tridosh also has impact on the health heart and blood vessels, and this physiological change in cardiovascular system is more prominent in the elderly population aged 60 and above.^[7]

In a recent study it was found that ageing is an important risk factor in the development of cardiovascular changes.^[5] Aging results in an increase in cardiovascular disease and a decrease in cardiac reserve at the same time that the repair processes designed to deal with these problems become less active/effective. 1) Structurally, the heart thickens and stiffens with age resulting in the increased imposition of a number of functional demands. 2) Functionally, aging causes significant functional deficits with exercise.^[4]

The relation between Doshas and Heart Rate Variability are highly intertwined with each other. Similarly, Heart Rate Variability and Nadi examination are also inter-related because the assessment of beat to beat variation is translated into the pulse examination.^[1] Thus, Nadi examination or radial pulse assessment and beat to beat variation may studied for exploring Dosha changes in an individual.^[1,12]

Heart Rate Variability (HRV) is variation in beat to beat intervals of the heart.^[6] Electrocardiographically, R wave represents the peak of QRS complex,^[13] and R to R duration is also called as RR interval; NN interval is used while assessing heart rate variability to highlight the fact that processed beats are normal beats.^[6]

HRV is an outcome of the impact of autonomic regulation on the heartbeat^[1] and Doshas are also autonomous regulators of body physiology.^[1] In recent years various scholars have extensively studied the impact of Doshas on HRV. Thus, Travis and Wallace also tried to explore the possible association between autonomous nervous system and Doshas.^[1,14]

Putting all these facts together, it may be inferred that ageing is linked with physiological changes in heart and vessels which is reflected by arterial stiffness. Similarly, the examination of pulse helps determine the Tridosha in individuals. Thus, an association may be proposed between Tridosha and pulse rate as well as RR interval which depict the physiological as well as pathological changes of the heart. Similarly, exercise has significant impact on the heart which is again measured by pulse rate. In conclusion, estimation heart rate variability through pulse rate and RR interval on lines of Dosha classification of elderly persons may yield crucial information about the individuals with maximum probability of having cardiovascular changes. Keeping this triangular conception, the present study was designed to measure the relation of Dosha-Prakruti in elderly using HRV.

METHODOLOGY

The present study was carried out in Aadhar Hospital located at Parwati Gaon, Pune under Department of Sharir Kriya, College of Ayurveda, Bharati Vidyapeeth, Pune, Maharashtra from January, 2019 to August, 2019. The trial was conducted after getting approval from ethical committee vide Letter No. BVDU/COA/46/2018-19. The voluntary written informed consent was sought from the trial participants before commencement of the trial. It was an observational study wherein participants were evaluated for HRV before and after exercise. The sample size was 30 participants aged 60 years and above. The intervention consisted of aerobic exercise i.e. cycling and walking for 60 minutes three times a week for 8 weeks with follow-up at every fortnight.

Inclusion criteria was participants in the age group of 45 years and above of both sexes and exclusion criteria consisted of participants suffering from any systemic disorder; especially

cardiac problems; pregnant and lactating women, and non-compliance of study protocol for one visit.

The study parameters included determination of Doshaj Prakruti using AyuSoft software in the target population,^[15] calculating heart rate variability (HRV) using RR interval of ECG,^[16] studying effect of exercise on HRV using Modified Baecke Questionnaire^[17] and effect of exercise on HRV using Lawton Broody Index^[18] in the study participants. The data on these parameters were collected before and after the trial.

RESULTS

The demographic data has given in Table 1.

Basic variables	No. of patients	%
Gender		
Female	13	43.3
Male	17	56.7
Total	30	100.0
Age in years		
<50	6	20.0
50-60	19	63.3
61-70	5	16.7
Total	30	100.0
Present illness		
NAD	21	70.0
Yes	9	30.0
• ARTHRITIS	3	10.0
• DM	2	6.7
• HTN	4	13.3

The distribution of Dosha has been shown in Table 2.

Table 2: Dosha distribution of participants studied.

	No. of patients (n=30)	%	Female (n=13)	%	Male (n=17)	%
Vata						
• 1-25	9	30.0	5	38.5	4	23.5
• 26-50	15	50.0	7	53.8	8	47.1
• 51-75	6	20.0	1	7.7	5	29.4
• 76-100	0	0.0	0	0.0	0	0.0
Pitta						
• 1-25	4	13.3	3	23.1	1	5.9
• 26-50	24	80.0	8	61.5	16	94.1
• 51-75	2	6.7	2	15.4	0	0.0
• 76-100	0	0.0	0	0.0	0	0.0

Kapha						
• 1-25	18	60.0	6	46.2	12	70.6
• 26-50	9	30.0	4	30.8	5	29.4
• 51-75	3	10.0	3	23.1	0	0.0
• 76-100	0	0.0	0	0.0	0	0.0

The study variables and their efficacy has been depicted in Tables 3, 4 and 5. Descriptive and inferential statistical analysis has been carried out in the present study presented on Mean±SD (Min-Max) and results on categorical measurements are presented in Number (%).

Table 3: Distribution of study variables.

Variables	Baseline	After study	% difference
R R interval			
<1	28(93.3%)	18(60%)	-33.3%
>1	2(6.7%)	12(40%)	33.3%
Total	30(100%)	30(100%)	-
Baecke Score			
1-5	9(30%)	8(26.7%)	-3.3%
5.1-10	21(70%)	22(73.3%)	3.3%
Total	30(100%)	30(100%)	-
Lawton Brody Score			
1	1(3.3%)	0(0%)	-3.3%
2	1(3.3%)	5(16.7%)	13.4%
3	6(20%)	5(16.7%)	-3.3%
4	9(30%)	9(30%)	0.0%
5 & more	13(43.3%)	11(36.7%)	-6.6%
Total	30(100%)	30(100%)	-

Significance is assessed at 5 % level of significance. Student t test (two tailed, independent) has been used to find the significance of study parameters on continuous scale between two groups (Inter group analysis) on metric parameters.

Table 4: Comparison of study variables at pre and post assessment.

	Before exercise	After Exercise	Difference	95% CI	t value	P value
Pulse rate	76.20±2.85	90.60±4.90	-14.400	16.234 to -12.566	16.063	<0.001**
R R interval	0.75±0.13	0.88±0.22	-0.130	-0.224 to -0.036	-2.840	0.008**
Baecke Score	5.91±1.48	5.74±1.73	0.168	-0.268 to 0.603	0.787	0.438
Lawton Brody Score	4.17±1.18	4.13±1.50	0.033	-0.388 to 0.455	0.162	0.873

Student *t* test

The Statistical software namely SPSS 22.0, and R environment ver.3.2.2 were used for the analysis of the data.

Variables	<50%	>50%	Total	P value
Vata				
Pulse rate				
Before Exercise	76.33±3.05	75.67±1.97	76.20±2.85	0.616
After Exercise	90.33±5.16	91.67±3.88	90.60±4.90	0.001**
R R interval				
Before Exercise	0.76±0.13	0.72±0.12	0.75±0.13	0.476
After Exercise	0.88±0.23	0.90±0.21	0.88±0.22	0.001**
Pitta				
Pulse rate				
Before Exercise	76.14±2.82	77.00±4.24	76.20±2.85	0.688
After Exercise	90.21±4.82	96±2.83	90.60±4.90	0.108
R R interval				
Before Exercise	0.75±0.13	0.80±0.14	0.75±0.13	0.568
After Exercise	0.87±0.22	1.05±0.21	0.88±0.22	0.263
Kapha				
Pulse rate				
Before Exercise	76.31±2.87	75.50±3.00	76.20±2.85	0.606
After Exercise	90.62±5.18	90.50±3.00	90.60±4.90	0.966
R R interval				
Before Exercise	0.76±0.13	0.70±0.08	0.75±0.13	0.401
After Exercise	0.90±0.22	0.73±0.10	0.88±0.22	0.130

DISCUSSION

The present study was an observational study to evaluate the effect of aerobic exercise on heart rate variability in various Dosha Prakruti. Before that, no trial has been conducted in the target population for establishing relation of Dosha with HRV. Thus, it is a naïve research step in correlation and contemporary interpretation of Dosha, Heart rate variability, and exercise in the designated population.

Regarding the age and gender, male constituted 57% and female were 43%. The data infers that male were more than the female. Consequently, the data may be extrapolated that male are more prone to develop the heart rate variability related problems as well due to ageing. The data is consistent with Andreas Voss *et al* who concluded that gender and age influences need to be considered when performing HRV studies.^[19] He found that males in the age group of 65 years and above had less heart rate as compared to female indicating that they may be compromised of cardiac problems in the coming days of their life. Similar, Jindal R *et al* reported that age has differential influence on vagal function in individuals in the age group of 60 years and above with implications for cardiovascular disease risk.^[20]

Moreover, the data also inferred that the mean age of the participants was 64 years in both the gender. Maximum number of participants was in the age group of 66 to 70 years (Table 1) which shows that with ageing the vagal stimulation is constantly compromised as evidenced by various models of ageing in humans.^[1,21]

Regarding Dosha distribution of study participants, the dominance of Vata, Pitta and Kapha in the elderly population was maximally 50%, 80% and 30% in the bracket of 1-25%, 26-50% and 26-50% respectively on AyuSoft proforma (Table 2). It inferred that overall; Pitta was dominant at the start of the trial. The data is not in conformity with that of Manohar et al who failed to show any Dosha imbalance in the study trial.^[1] Similarly, in the evaluation of data on exercise in relation to HRT, Vata Dosha was found dominant. More prospective trials are needed to explore the Dosha imbalance in this regard.

Regarding present illness of the participants, It was observed that majority (70%) of the participants were free from any co-morbid conditions except 6 cases who had co-morbid conditions of DM and hypertension (Table 1). The data seems compromised as the participants with co-morbid conditions are the prime source of confounding and bias. Hence, it is primarily required to select a homogenous population free from confounders. Besides this, Manohar et al has tried to show the HRV and its relation with Dosha in various morbid conditions^[1] but the study is miniature in condition as the single disease or health status focused prospective studies are need of the hour in Ayurveda.

Regarding pulse rate, there was an increment in pulse rate of 53.3% as compared to the baseline data after exercise. Maximum increase in pulse rate was noticed in the bracket of 90-99 beats as compared to 71-79 (Table 3). Aerobic exercise has multiple positive effects on health outcomes in healthy elderly individuals including increased cardiovascular performance^[1,22] Contrary to this, Hartaigh *et al* found that twelve months of moderate intensity aerobic training did not improve RPR among older adults.^[23] Hence, more prospective trials are needed to prove the same.

Regarding RR interval, there was constant improvement as 93.3% (28) had RR interval <1 at baseline which increased to 60% (18) had it >1 showing the strong indication of cardiac output (Table 3). The data is in constant with Manohar et al.^[1] In a major prospective population based study, it was observed that modest aerobic exercise throughout the aging process is potentially related with minimum ECG changes.^[24]

Thus putting altogether, heart diseases such as arrhythmias are prevalent in aging population and are linked to increased electrophysiological and autonomic dysfunction in the heart.^[25] Heart rate variability (HRV) is the indicator of cardiac electrophysiology and autonomic regulation including respiratory, baroreflex, and circadian fluctuations indicative of healthier heart.^[26] In the same context, increased HRV may exhibit abnormal sinus patterns with advancing age,^[27] but physical activity in the middle age is associated with more favorable HRV data which reflects the increased vagal modulation and reduced sympathetic activity of the heart.^[28]

Estimation of the Dosha imbalance is highly importance in Ayurvedic diagnosis and treatment of diseases. But the assessment is primarily subjective in nature with no valid tool. Examination of pulse is an important tool of Ayurvedic medicine used to seek the insights into Dosha with higher inter-variability.

The present trial has been focused on interpretation of Dosha imbalance and determination using sophisticated tools such as AyuSoft proforma to overcome the inter-variability. On the other hand, clinical determination of Dosha imbalance is done through pulse examination which is again at the risk of inter-variability. Thus, heart rate variability has also served an indirect method of Dosha imbalance in the target population. For that, series of studies comparing HRV spectral analysis with clinical assessment of Dosha dynamics can predict doshic imbalance.

CONCLUSION

The clinical implication of this study is that recommending exercise in Vata Dosha individuals may better the heart health in the older adults and elderly. The limitations of the study are small sample size and single arm study. Hence, it is recommended for the future researchers to design a randomized controlled trial to establish the relation of HRV to Dosha in more scientific way.

ACKNOWLEDGEMENT

Authors are highly grateful to the Vice-chancellor and Dean, faculty of Ayurveda for their generous support in accomplishment of this valuable research work.

REFERENCES

1. Ram Manohar P, Sorokin O, Chacko J, Nampoothiri V. An exploratory clinical study to determine the utility of heart rate variability analysis in the assessment of dosha imbalance. *J Ayurveda Integr Med [Internet]*, 2018; 9(2): 126–30.
2. Narayanaswamy V. Origin and development of ayurveda: (a brief history). *Anc Sci Life [Internet]*, 1981; 1(1): 1–7.
3. Closshey JE. The Ayurveda Tridosha concept and the human apolipoprotein genotype: A quatuatative study, 2015.
4. Fajemiroye JO, Cunha LC Da, Saavedra-Rodríguez R, Rodrigues KL, Naves LM, Mourão AA, et al. Aging-Induced Biological Changes and Cardiovascular Diseases. *Biomed Res Int [Internet]*, 2018; 10: 1–14.
5. Steenman M, Lande G. Cardiac aging and heart disease in humans. *Biophys Rev [Internet]*, 2017; 20, 9(2): 131–7.
6. Harupjit Singh. Relationship between Three Ayurvedic Doshas and Heart Rate Variability Frequency Bands: A Pilot Study. Thapar University, Patiala, 2016.
7. Vagbhata. *Ashtanga Hridayam [Internet]*. 1st ed. R. Vidyanath, editor. Varansi: Chaukhamba Surbharati Prakashan, 2013; 2–10, 12–32, 45–49.
8. Lad V. *Textbook of Ayurveda. Vol.1. 1st ed.* Albuquerque, New Mexico: Ayurvedic Press, 2002; 1–2.
9. Jaiswal YS, Williams LL. A glimpse of Ayurveda – The forgotten history and principles of Indian traditional medicine. *J Tradit Complement Med [Internet]*, 2017 1; 7(1): 50–3.
10. World Health Organization (WHO). Benchmarks for training in traditional/ complementary and alternative medicine Benchmarks for Training in Ayurveda. Geneva: World Health Organization, 2010; 23–25, 33 - 41.
11. Kumar Meena D, Upadhyay D, Singh R, Dwibedy BK, ResidentIII J, Resident S, et al. A critical review of fundametal principales of Ayurveda. *Int Ayurvedic Med J [Internet]*, 2015; 3(7): 2075–80.
12. Pradnya D, Nalini A, Nagarathna R, Raju T, Sendhilkumar R, Meghana A, et al. Effect of yoga as an add-on therapy in the modulation of heart rate variability in children with duchenne muscular dystrophy. *Int J Yoga*, 2019; 12(1): 55.
13. European Society of Cardiology and the North American Society of Pacing and Electrophysiology. Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation [Internet]*, 1996; 1,

- 93(5): 1043–65.
14. Travis F, Wallace R. Dosha brain-types: A neural model of individual differences. *J Ayurveda Integr Med [Internet]*, 2015; 1, 6(4): 280.
 15. Ghate US, Indapurkar K V. Study of Dhatu Sarata and its association with blood indices. *International J Res Ayurveda Pharm [Internet]*, 2014; 4, 5(3): 309–14.
 16. Hansen AL, Johnsen BH, Thayer JF. Relationship between heart rate variability and cognitive function during threat of shock. *Anxiety, Stress Coping [Internet]*, 2009; 22(1): 77–89.
 17. Hertogh EM, Monninkhof EM, Schouten EG, Peeters PH, Schuit AJ. Validity of the Modified Baecke Questionnaire: comparison with energy expenditure according to the doubly labeled water method. *Int J Behav Nutr Phys Act [Internet]*, 2008; 5(1): 30.
 18. De Meersman RE. Heart rate variability and aerobic fitness. *Am Heart J [Internet]*, 1993; 125(3): 726–31.
 19. Voss A, Schroeder R, Heitmann A, Peters A, Perz S. Short-Term Heart Rate Variability—Influence of Gender and Age in Healthy Subjects. Hernandez A V, editor. *PLoS One [Internet]*, 2015; 30, 10(3): e0118308.
 20. Jindal RD, Vasko RC, Jennings JR, Fasiczka AL, Thase ME, Reynolds CF. Heart Rate Variability in Depressed Elderly. *Am J Geriatr Psychiatry [Internet]*, 2008; 16(11): 861–6.
 21. Zhang J. Effect of Age and Sex on Heart Rate Variability in Healthy Subjects. *J Manipulative Physiol Ther [Internet]*, 2007; 30(5): 374–9.
 22. Vigorito C, Giallauria F. Effects of exercise on cardiovascular performance in the elderly. *Front Physiol [Internet]*, 2014; 5(51): 1–8.
 23. O'Hartaigh B, Pahor M, Buford TW, Dodson JA, Forman DE, Gill TM. Physical activity and resting pulse rate in older adults: Findings from a randomized controlled trial. *Am Heart J [Internet]*, 2014; 168(4): 597–604.
 24. Soares-Miranda L, Sattelmair J, Chaves P, Duncan GE, Siscovick DS, Stein PK, et al. Physical Activity and Heart Rate Variability in Older Adults. *Circulation [Internet]*, 2014; 27, 129(21): 2100–10.
 25. Tsuji H, Venditti FJ, Manders ES, Evans JC, Larson MG, Feldman CL, et al. Reduced heart rate variability and mortality risk in an elderly cohort. The Framingham Heart Study. *Circulation [Internet]*, 1994; 90(2): 878–83.
 26. Task Force of the European Society of Cardiology the North American Society of Pacing Electrophysiology. Standards of measurement, physiological interpretation, and clinical

- use. *Circulation*, 1996; 93(5): 1043–65.
27. Kleiger RE, Stein PK, Bigger JT. Heart Rate Variability: Measurement and Clinical Utility. *Ann Noninvasive Electrocardiol* [Internet], 2005; 10(1): 88–101.
28. Hull SS, Vanoli E, Adamson PB, Verrier RL, Foreman RD, Schwartz PJ. Exercise training confers anticipatory protection from sudden death during acute myocardial ischemia. *Circulation* [Internet], 1994; 89(2): 548–52.

WJPPS COPY PROOF